



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

otropically) in darkness, and give a very characteristic geotropic reaction, having abundant statolith starch. *P. epiphylla* generally does not grow in darkness. Neither these two nor *Lophocolea bidentata* are geotropic in light. The behavior in darkness of *P. epiphylla*, the two species of *Lophocolea*, and the sporogone of *Aneura* must be considered as purposeless.

The researches appear to strengthen the statolith theory of geotropic perception.—C. R. B.

Photosynthesis.—USHER and PRIESTLY, having shown in a previous paper¹⁵ that CO_2 may be decomposed in plants independently of enzymic or "vital" activity provided the products (H_2O_2 and $\text{H}'\text{COH}$) are removed, have now devised an arrangement by which photolysis of CO_2 can be produced *in vitro*.¹⁶ By covering gelatin plates, or even water, with a very thin uniform layer of chlorophyll deposited from solution, and placing the glass plate or the shallow dish of water carrying the film in a vessel with CO_2 and illuminating it, they found formaldehyde produced and were able to recover identifiable quantities. By making up the gelatin with an aqueous solution of a catalase, the hydrogen peroxid also produced was decomposed with the evolution of O_2 , 2^{cc} being obtained in one case. The chlorophyll was gradually bleached, in harmony with the view that it acts as a sensitizer and is destroyed in the process of photolysis. Synthesis of $\text{H}'\text{COH}$ into carbohydrate was found in the earlier paper to be dependent on the healthy condition of the protoplast, and feeding experiments have already shown that $\text{H}'\text{COH}$ when supplied in very dilute form can be condensed by green plants if illuminated. The authors painted the white petals of *Saxifraga Wallacei* with chlorophyll, and floated them on water charged with CO_2 in light. In the course of a day they were found to contain starch. Thus they secured photosynthesis by a different (though inefficient) arrangement of the mechanism of a green leaf.

Further study was made of photolysis of CO_2 in the presence of uranium salts. They have direct evidence of the production of formic acid (which also is produced under some conditions in the plant as an intermediate product in the reduction of CO_2), but they were unable to isolate and identify formaldehyde.

These papers record a most important step in solving the problems of photosynthesis.—C. R. B.

Vascular anatomy of cycads.—Two years ago MATTE published his thesis on the vascular anatomy of the cycads,¹⁷ in which he presented in great detail, with copious illustrations, the vascular anatomy of the leaves and flowers in representative species of all the nine genera; and of the seedlings of *Dioon edule*, *Cycas*

¹⁵ USHER, F. L., and PRIESTLY, J. H., A study of the mechanism of carbon assimilation in green plants. Proc. Roy. Soc. London B. 77:369-376. 1905.

¹⁶ —The mechanism of carbon assimilation in green plants: the photolytic decomposition of carbon dioxid *in vitro*. *Idem* 78:318-327. 1906.

¹⁷ MATTE, H., Recherches sur l'appareil libero-ligneux des Cycadées. pp. 233. pls. 16. figs. 264. 1904.